

ON A STRATIFICATION OF THE MODULI $\overline{M}_{0,n}(G(2,4), d)$ AND ENUMERATIVE GEOMETRY.

CRISTINA MARTÍNEZ RAMIREZ

ABSTRACT. We study the geometry of the moduli space $\overline{M}_{0,n}(G(2,4), d)$. We consider a particular stratification of this moduli space. As an application we compute the degree of the variety parametrizing rational ruled surfaces with a minimal directrix of degree $\frac{d}{2} - 1$ by studying divisors in this moduli space of stable maps.

1. INTRODUCTION

The geometry of the Kontsevich's moduli space $\overline{M}_{g,n}(\mathbb{P}^r, d)$ of degree d stable maps from n -pointed, genus g curves to $\mathbb{P}^r$ has been studied intensively in the literature. R. Vakil studied its connection with the enumerative geometry of rational and elliptic curves in projective space in [Vak]. R. Pandharipande studied the theory of $\mathbb{Q}$ -Cartier divisors on this space in [Pan1] and proved an algorithm for computing all characteristic numbers of rational curves in $\mathbb{P}^r$. R. Pandharipande also computed the degree of the 1-cuspidal rational locus in the linear system of degree d plane curves.

The enumerative geometry of rational ruled surfaces of degree d in $\mathbb{P}^3$ is closely related to the intrinsic geometry of the moduli space $\overline{M}_{0,n}(G(2,4), d)$, of Kontsevich stable maps from n -pointed genus 0 curves to the Grassmannian $G(2,4)$ of lines in $\mathbb{P}^3$, representing d times the positive generator of the homology group $H_2(G(2,4), \mathbb{Z})$, (see [Mar]).

In [Mar] we solved the enumerative problem of computing the degree of the Severi variety of degree d rational ruled surfaces in the ambient projective space of surfaces of degree d in $\mathbb{P}^3$. Here we study the more refined problem of enumerating rational ruled surfaces with a fixed directrix of minimum degree. In particular we compute the degree of the codimension one subvariety of rational ruled surfaces with a minimal directrix of degree $\frac{d}{2} - 1$ by intersecting divisors in the moduli space of stable maps $\overline{M}_{0,0}(G(2,4), d)$.

In a first step, we use the variety R_d^0 of degree d morphisms from $\mathbb{P}^1$ to the Grassmannian $G(2,4)$, as a parameter space for parametrized rational ruled surfaces in $\mathbb{P}^3$. We consider a certain stratification of the variety R_d^0 defined as follows. If $\mathcal{Q}$ is the universal bundle on $G(2,4)$, the stratum is defined by the following locally closed condition:

$$R_{d,a}^0 = \{f \in R_d^0 \mid \dim H^1(\mathbb{P}^1, f^* \mathcal{Q}^\vee \otimes \mathcal{O}_{\mathbb{P}^1}(-a-2)) \geq 1 \text{ and} \\ \dim H^0(\mathbb{P}^1, f_q^* \mathcal{Q}^\vee \otimes \mathcal{O}_{\mathbb{P}^1}(-a-1)) = 0\}.$$

2000 *Mathematics Subject Classification.* Primary 14N35, 14N10; Secondary 14C05, 14C15.

Key words and phrases. Enumerative geometry, intersection theory, stable maps.

This has a geometrical interpretation as rational ruled surfaces with a minimal directrix of degree a . First, we consider the closure $R_{d,a}$ of the stratum $R_{d,a}^0$ in the Quot scheme compactification R_d of the space of morphisms R_d^0 . We prove that the subschemes $R_{d,a}$ are irreducible and have the expected codimension. We give an explicit formula for their Poincaré duals derived from Porteous formula. In the last section, we consider their closure in the Kontsevich compactification of stable maps $\overline{M}_{0,3}(G(2,4), d)$. We show that the given stratification of the space R_d^0 can be extended to $\overline{M}_{0,3}(G(2,4), d)$, but in this case the codimension of the stratum is different, that is, not all boundary points correspond to closure points of the open set $R_{d,a}^0$, except for the big stratum of rational ruled surfaces with a minimal directrix of degree $\frac{d}{2} - 1$. In this case we compute its degree by interpreting it as a tautological intersection on the space of stable maps.

1.1. Conventions. We work over the field of complex numbers $\mathbb{C}$. By a scheme we mean a scheme of finite type over $\mathbb{C}$. By a variety, we mean a separated integral scheme. Curves are assumed to be complete and reduced. For a variety X , $A_d X$ and $A^d X$ can be taken to be the Chow homology and cohomology groups. $A^d X$ and $A_{n-d} X$ are identified by the Poincaré duality isomorphism. For $\beta \in A_k X$, $\int_\beta c$ is the degree of the zero cycle obtained by evaluating c_k on β , where c_k is the component of c in $A^k X$. If $\mathcal{F}$ is a sheaf over a curve C , $h^0(C, \mathcal{F})$ and $h^1(C, \mathcal{F})$ denote the dimensions of the cohomology groups $H^0(C, \mathcal{F})$ and $H^1(C, \mathcal{F})$.

For any formal series $a(t) = \sum_{k=-\infty}^{+\infty} a_k t^k$, we set

$$\Delta_{p,q}(a) = \det \begin{pmatrix} a_p & \cdots & a_{p+q-1} \\ \vdots & & \vdots \\ a_{p-q+1} & \cdots & a_p \end{pmatrix}.$$

2. A STRATIFICATION OF THE SPACE R_d^0 AND ENUMERATIVE GEOMETRY

Let R_d^0 be the variety of degree d morphisms from $\mathbb{P}^1$ to $G(2,4)$. We will denote R_d^{00} the Zariski open set in R_d^0 corresponding to a morphism birational onto its image. The Grothendieck Quot scheme R_d parametrizing rank 2 and degree d quotients of a trivial vector bundle $\mathcal{O}_{\mathbb{P}^1}^4$, is a natural projective scheme compactifying R_d^0 . We will call it R_d . We consider the evaluation map from $R_d^0 \times \mathbb{P}^1$ to the Grassmannian $G(2,4)$, and the projection map over the first component.

$$(1) \quad \begin{array}{ccc} R_d^0 \times \mathbb{P}^1 & \xrightarrow{e} & G(2,4) \\ \pi_1 \downarrow & & \\ R_d^0 & & \end{array}$$

The pull back of the universal exact sequence on $G(2,4)$, gives a universal exact sequence on R_d^0 which is the restriction of the universal quotient over $R_d \times \mathbb{P}^1$.

For $f \in R_d^0$, the pull-back under f of the universal quotient on the Grassmannian $f^* \mathcal{Q}$, is a bundle of rank 2 on $\mathbb{P}^1$ isomorphic to $\mathcal{O}_{\mathbb{P}^1}(a) \oplus \mathcal{O}_{\mathbb{P}^1}(d-a)$

where $0 \leq a \leq \frac{d}{2}$, in particular $a \leq d-a$. There is a canonical exact sequence

$$0 \rightarrow N \rightarrow \mathcal{O}_{\mathbb{P}^1}^4 \rightarrow \mathcal{O}(a) \oplus \mathcal{O}(d-a) \rightarrow 0$$

where N is a rank 2 and degree $-d$ bundle. Thus, we get a morphism $F : \mathbb{P}(\mathcal{O}(a) \oplus \mathcal{O}(d-a)) \rightarrow \mathbb{P}^3$. The image of this morphism is a *scroll*. We will denote it by $X_{a,d-a}$.

Definition 2.1. For each $0 \leq a \leq \frac{d}{2}$, $R_{d,a}^0$ is the subvariety of R_d^0 consisting of those $f \in R_d^0$ such that $f^* \mathcal{Q} \cong \mathcal{O}_{\mathbb{P}^1}(a) \oplus \mathcal{O}_{\mathbb{P}^1}(d-a)$.

Proposition 2.2. $R_{d,a}^0$ is locally closed in R_d^0 , irreducible and it has dimension $4d+4$ if $d=2a$, and $3d+5+2a$, if $d > 2a$.

Proof. The set $R_{d,a}^0$ is locally closed by [Sha].

Let $F_a \subset \mathbb{P} \operatorname{Hom}(\mathcal{O}^4, \mathcal{O}_{\mathbb{P}^1}(a) \oplus \mathcal{O}_{\mathbb{P}^1}(d-a))$ be the open set of epimorphisms

$$\mathcal{O}^4 \twoheadrightarrow \mathcal{O}_{\mathbb{P}^1}(a) \oplus \mathcal{O}_{\mathbb{P}^1}(d-a).$$

F_a is in bijection with the set of locally free rank 2 quotients of $\mathcal{O}_{\mathbb{P}^1}^4$ isomorphic to $\mathcal{O}_{\mathbb{P}^1}(a) \oplus \mathcal{O}_{\mathbb{P}^1}(d-a)$ modulo $\mathbb{C}^*$. Let $F_a \xrightarrow{\psi} R_{d,a}^0$ be the surjective morphism such that the image by ψ of each quotient is its class of isomorphism in $R_{d,a}^0$. The fibers are isomorphic to $\mathbb{P}(\operatorname{Aut}(\mathcal{O}_{\mathbb{P}^1}(a) \oplus \mathcal{O}_{\mathbb{P}^1}(d-a)))$.

A dimension calculation yields:

(1) If $a = d-a$,

$$\dim \mathbb{P}(\operatorname{Aut}(\mathcal{O}(a) \oplus \mathcal{O}(d-a))) = 3.$$

(2) If $d-a > a$,

$$\dim \mathbb{P}(\operatorname{Aut}(\mathcal{O}(a) \oplus \mathcal{O}(d-a))) = d-2a+2.$$

The dimension of F_a is given by

$$\dim \mathbb{P} \operatorname{Hom}(\mathcal{O}_{\mathbb{P}^1}^4, \mathcal{O}_{\mathbb{P}^1}(a) \oplus \mathcal{O}_{\mathbb{P}^1}(d-a)) = \dim F_a = 4d+7.$$

By the fiber dimension theorem ([Sha], §I.6.3),

$$(2) \quad \dim R_{d,a}^0 = \begin{cases} 4d+4 & \text{if } d=2a \\ 3d+5+2a & \text{if } d>2a \end{cases}$$

In particular this proves that the subvarieties $R_{d,a}^0$ are irreducible, because F_a is irreducible. $\square$

Lemma 2.3. $q \in R_{d,a}^0$ if and only if $h^0(\mathbb{P}^1, f_q^* \mathcal{Q}^\vee \otimes \mathcal{O}_{\mathbb{P}^1}(a)) \geq 1$ and

$$h^0(\mathbb{P}^1, f_q^* \mathcal{Q}^\vee \otimes \mathcal{O}_{\mathbb{P}^1}(a-1)) = 0.$$

Proof. Let q be a point in $R_{d,a}^0$, and $f_q : \mathbb{P}^1 \rightarrow G(2,4)$ the morphism corresponding to that point. Suppose $f_q^* \mathcal{Q}$ decomposes as $\mathcal{O}_{\mathbb{P}^1}(a) \oplus \mathcal{O}_{\mathbb{P}^1}(d-a)$. Tensoring with the linear bundle $\mathcal{O}_{\mathbb{P}^1}(a)$,

$$f_q^* \mathcal{Q}^\vee \otimes \mathcal{O}_{\mathbb{P}^1}(a) = \mathcal{O}_{\mathbb{P}^1}(2a-d) \oplus \mathcal{O}_{\mathbb{P}^1},$$

and tensoring with $\mathcal{O}_{\mathbb{P}^1}(a-1)$, we have

$$f_q^* \mathcal{Q}^\vee \otimes \mathcal{O}_{\mathbb{P}^1}(a-1) = \mathcal{O}_{\mathbb{P}^1}(2a-d-1) \oplus \mathcal{O}_{\mathbb{P}^1}(-1).$$

Since $a \leq d-a$, it follows that $h^0(\mathbb{P}^1, f_q^* \mathcal{Q}^\vee \otimes \mathcal{O}_{\mathbb{P}^1}(a)) \geq 1$, and

$$h^0(\mathbb{P}^1, f_q^* \mathcal{Q}^\vee \otimes \mathcal{O}_{\mathbb{P}^1}(a-1)) = 0.$$

Conversely, if $f_q^* \mathcal{Q}^\vee \cong \mathcal{O}(-n) \oplus \mathcal{O}_{\mathbb{P}^1}(n-d)$ with $n \leq d-n$, then

$$f_q^* \mathcal{Q}^\vee \otimes \mathcal{O}_{\mathbb{P}^1}(a) = \mathcal{O}_{\mathbb{P}^1}(n-d+a) \oplus \mathcal{O}_{\mathbb{P}^1}(a-n),$$

then $h^0(\mathbb{P}^1, f_q^* \mathcal{Q}^\vee \otimes \mathcal{O}_{\mathbb{P}^1}(a)) \geq 1$ implies $a \geq n$ and

$h^0(\mathbb{P}^1, f_q^* \mathcal{Q}^\vee \otimes \mathcal{O}_{\mathbb{P}^1}(a-1)) = 0$ implies $a \leq n$, therefore $n = a$ and $q \in R_{d,a}^0$.
□

Remark 1. We observe that the cases $d = 2a$ and $d = 2a + 1$ are special. In these cases $R_{d,a}^0$ has the maximum dimension. This means that $R_{d,a}^0$ is an open set in R_d^0 and therefore the corresponding degree has already been computed in ([Mar]).

Remark 2. By the Serre duality theorem and Lemma 2.3 the subvarieties $R_{d,a}^0$ can also be defined as

$$(3) \quad R_{d,a}^0 = \{f \in R_d^0 \mid h^1(\mathbb{P}^1, f^* \mathcal{Q} \otimes \mathcal{O}_{\mathbb{P}^1}(-a-2)) \geq 1 \text{ and } h^1(\mathbb{P}^1, f^* \mathcal{Q} \otimes \mathcal{O}_{\mathbb{P}^1}(-(a-1)-2)) = 0\}.$$

Note that when $d > 2a$ a slightly stronger result is true:

$q \in R_{d,a}^0$ if and only if

$$h^0(\mathbb{P}^1, f_q^* \mathcal{Q}^\vee \otimes \mathcal{O}_{\mathbb{P}^1}(a)) = 1 \text{ and } h^0(\mathbb{P}^1, f_q^* \mathcal{Q}^\vee \otimes \mathcal{O}_{\mathbb{P}^1}(a-1)) = 0.$$

These sets are locally closed by the Semicontinuity Theorem.

Lemma 2.4. *The Zariski closure $\overline{R}_{d,a}^0$ of the stratum $R_{d,a}^0$ in R_d^0 coincides with*

$$(4) \quad \{f \in R_d^0 \mid h^1(\mathbb{P}^1, f^* \mathcal{Q} \otimes \mathcal{O}_{\mathbb{P}^1}(-a-2)) \geq 1\}.$$

Proof. By definition, the closure $\overline{R}_{d,a}^0$ is the minimum closed set containing $R_{d,a}^0$. We observe that $R_{d,a}^0$ is contained in the set given in (4) that it is closed, therefore it also contains $\overline{R}_{d,a}^0$. Now we notice that every closed set containing $R_{d,a}^0$ contains the closed set

$$\{f \in R_d^0 \mid h^0(\mathbb{P}^1, f_q^* \mathcal{Q}^\vee \otimes \mathcal{O}_{\mathbb{P}^1}(a)) \geq 1, \text{ and } h^0(\mathbb{P}^1, f_q^* \mathcal{Q}^\vee \otimes \mathcal{O}_{\mathbb{P}^1}(a-1)) \geq 0\},$$

but the second closed condition is satisfied trivially, therefore the closure $\overline{R}_{d,a}^0$ coincides with the set (4).
□

Remark 3. We observe that if $f_q^* \mathcal{Q} \cong \mathcal{O}_{\mathbb{P}^1}(-n) \oplus \mathcal{O}_{\mathbb{P}^1}(n-d)$ with $n \leq d-n$, $h^0(\mathbb{P}^1, f_q^* \mathcal{Q}^\vee \otimes \mathcal{O}_{\mathbb{P}^1}(a)) \geq 1$ implies $a \geq n$ and therefore $\overline{R}_{d,a}^0 = R_{d,a}^0 \cup R_{d,a-1}^0 \cup \dots$

Lemma 2.5. *The subvariety $R_{d,a}^0$ parametrizes rational ruled surfaces with a minimal directrix of degree a .*

Proof. The projection

$$\pi_a : \mathcal{O}_{\mathbb{P}^1}(a) \oplus \mathcal{O}_{\mathbb{P}^1}(d-a) \rightarrow \mathcal{O}_{\mathbb{P}^1}(a),$$

gives a section $\mathbb{P}(\mathcal{O}_{\mathbb{P}^1}(a)) \hookrightarrow \mathbb{P}(f^* \mathcal{Q})$. This section is mapped to a rational curve C^a of degree a . A study of the map

$$(5) \quad \mathcal{O}_{\mathbb{P}^1}(a) \oplus \mathcal{O}_{\mathbb{P}^1}(d-a) \rightarrow \mathcal{O}_{\mathbb{P}^1}(k),$$

shows that the map is surjective for $k \geq a$, therefore C^a is a *directrix* of minimal degree and (3) has a geometrical interpretation as the rational ruled surfaces with a minimal directrix of degree a . $\square$

3. CHANGE OF BASIS IN $Pic(R_d)$.

In this section we describe geometric generators for the Picard group of the Quot scheme R_d .

We consider the cohomology ring of the Grassmannian $G(2,4)$ and the universal sequence over it:

$$(6) \quad 0 \rightarrow \mathcal{N} \rightarrow \mathcal{O}_{G(2,4)}^4 \rightarrow \mathcal{Q} \rightarrow 0.$$

The special Schubert cycles on $G(2,4)$ are be represented as Chern classes of the universal quotient bundle and the universal subbundle:

$$T_1 = c_1(\mathcal{Q}) = c_1(\mathcal{N}^\vee), \quad T_b = c_2(\mathcal{N}), \quad T_a = c_2(\mathcal{Q}).$$

Also, $T_3 \in H_2(G(2,4), \mathbb{Z})$ will stand for the class of a line and T_4 for the class of a point in $G(2,4)$. The Poincare dual of the hyperplane class $c_1(\mathcal{N}^\vee)$ in $H^2(G(2,4), \mathbb{Z})$ determines the Plücker embedding of the Grassmannian $G(2,4)$ as a quadric in $\mathbb{P}^5$, that corresponds to a variety of lines in $\mathbb{P}^3$. Then the cycles defined above, have an interpretation as lines in $\mathbb{P}^3$. The cycle T_1 corresponds to lines in $\mathbb{P}^3$ meeting a given line, T_b corresponds to lines contained in a given plane and T_a to lines containing a given point. determines the Plücker embedding.

Let $e : R_d^0 \times \mathbb{P}^1 \rightarrow G(2,4)$ be the evaluation map. The following sets of morphisms define Weil divisors on R_d^0 :

- (A) The locus of morphisms whose image meets an a -plane T_a in the Grassmannian associated to a point $P \in \mathbb{P}^3$. We denote it by:

$$D := \{\varphi \in R_d^0 \mid e(t, \varphi) \cap T_a \neq \emptyset\}.$$

- (B) The set of morphisms Y sending a fixed point $t \in \mathbb{P}^1$ to a hyperplane T_1 on the Grassmannian:

$$Y := \{\varphi \in R_d^0 \mid e(t, \varphi) \in T_1 \text{ for a fixed } t \in \mathbb{P}^1\}.$$

Since the boundary of R_d is of codimension greater than 1 (see [Ber2]), these divisors extend to divisors in R_d , that is $A^1(R_d^0) = A^1(R_d)$. Let Y, D also denote the divisors on R_d . It is clear by the definition that when we move the point $t \in \mathbb{P}^1$ we get a linearly equivalent divisor. If we index the points t_i in $\mathbb{P}^1$, Y_i will denote the corresponding associated divisor. In the same way, every a -plane T_a in $G(2,4)$ is associated to a point P in $\mathbb{P}^3$ and if we move the point we also get an equivalent divisor that can be indexed as well. The divisors Y_i were already considered by Bertram in [Ber1] where he proved a moving lemma stating that these varieties can be made to intersect transversally. S.A. Stromme in [Str] gives a basis for the Picard group $A^1(R_d)$, formed by the divisors:

$$\begin{aligned} \alpha &= c_1(\pi_{1*}(\mathcal{E} \otimes \pi_2^* \mathcal{O}_{\mathbb{P}^1}(d)) - c_1(\pi_{1*}(\mathcal{E} \otimes \pi_2^* \mathcal{O}_{\mathbb{P}^1}(d-1))), \\ \beta &= c_1(\pi_{1*}(\mathcal{E} \otimes \pi_2^* \mathcal{O}_{\mathbb{P}^1}(d-1)), \end{aligned}$$

where $\mathcal{E}$ is the universal quotient over $R_d \times \mathbb{P}^1$, and π_1, π_2 are the projections maps over the first and second factors respectively.

Let h be the positive generator of the Picard group of $\mathbb{P}^1$ as well as its pull-back to $R_d \times \mathbb{P}^1$.

The Chern classes of the universal subbundle $\mathcal{K}$ are described in the proof of theorem 5.3 of [Str], that is

$$A(R_d \times \mathbb{P}^1) = \{A(R_d)[h], \mid h^2 = 0\}$$

Every class $z \in A(R_d \times \mathbb{P}^1)$ can be written in the form $z = x + h y$, with $y = \pi_{1*}(h) \in A(R_d)$, and $x = \pi_{1*}(h z) \in A(R_d)$,

$$(7) \quad c_i(\mathcal{K}) = t_i + h u_{i-1} \quad (1 \leq i \leq 2),$$

$t_i \in A^i(R_d)$, $u_i \in A^{i-1}(R_d)$, $i = 1, 2$ as in [Str], in particular $u_0 = -d$.

We express the classes t_1, u_1 in terms of the generators α and β .

Lemma 3.1. *The change of basis is*

$$\begin{aligned} \alpha &= -t_1, \\ \beta &= u_1. \end{aligned}$$

Proof. We need to compute the first Chern class of the bundle,

$$B_m = \pi_{1*}(\mathcal{E} \otimes \pi_2^* \mathcal{O}_{\mathbb{P}^1}(m)).$$

By Riemman Roch, we have that

$$\text{ch}(B_m) = \pi_{1*}((1 + h) \cdot \text{ch}(\mathcal{E} \otimes \pi_2^* \mathcal{O}_{\mathbb{P}^1}(m))).$$

The Chern classes of $\mathcal{E}$ can be computed from those of $\mathcal{K}$:

$$\begin{aligned} c_1(\mathcal{E}) &= -c_1(\mathcal{K}) = -t_1 + d h, \\ c_2(\mathcal{E}) &= c_1^2(\mathcal{K}) - c_2(\mathcal{K}) = t_1^2 - 2d h t_1 - t_2 - u_1 h. \end{aligned}$$

It follows that,

$$c_1(B_m) = (d - 1 - m) t_1 + u_1.$$

□

Lemma 3.2. *The change of basis from the α, β divisors to the geometric divisors defined above is:*

$$\begin{aligned} \alpha &= Y, \\ \beta &= -2dY + D. \end{aligned}$$

Proof. By the universal property of the Grassmannian, the pull back under the evaluation map e of the universal exact sequence (6) on $G(2, 4)$, gives us a universal exact sequence on $R_d^0 \times \mathbb{P}^1$ which is the restriction of the universal one on $R_d \times \mathbb{P}^1$. Therefore, by the description of the special Schubert cycles given, the class $[Y]$ of the divisor Y in the Chow ring $A(R_d)$ can be identified with the pushforward of the projection map over the first factor $\pi_{1*}(he^*(T_1)) = \pi_{1*}(e^*(c_1(\mathcal{Q})) \cdot h) = \pi_{1*}(h c_1(\mathcal{E}))$, where $\mathcal{E}$ is the universal quotient over $R_d \times \mathbb{P}^1$. Similarly, the class of the divisor D can be identified with $\pi_{1*}(c_2(\mathcal{E}))$.

The Riemann-Roch-Grothendieck theorem shows that the geometric divisor Y is just the divisor α . Now, we see that $[D] = \pi_{1*}(t_1^2 - 2t_1 d h - t_2 + h u_1) =$

$-2dt_1 + u_1 = 2d\alpha + \beta$, where β is the other generator of the Picard group of R_d . In particular, this also proves that the divisors D and Y constitute a basis for the Picard group $A^1(R_d)$. $\square$

Let P_d be the degree of R_d by the morphism induced by the hyperplane class in R_d . By lemma 3.2, P_d is the degree of the top codimensional cohomology class given by the selfintersection,

$$(8) \quad P_d = \int_{[R_d]} [Y]^{4d+4} \cap [R_d] = \int_{[R_d]} \alpha^{4d+4} \cap [R_d].$$

This intersection number is computed in [RRW] via Quantum cohomology. It is a Gromov-Witten invariant and can be obtained too by using the formulas of Vafa and Intriligator, [Ber1].

In [Mar] we prove that the intersection $Y_1 \cap Y_2 \cap Y_3 \cap D_1 \cap \dots \cap D_{4d+1}$ has no enumerative meaning in R_d and we compute it by reinterpreting it as a Gromov-Witten invariant in the Kontsevich moduli space of stable maps $\overline{M}_{0,4d+4}(G(2,4), d)$. For $d \geq 3$, this intersection is $d^3 \cdot Q_d$, where Q_d is the Severi degree of the variety of rational ruled surfaces in $\mathbb{P}^3$, and the contribution factor d^3 comes from the multiple covers.

4. CLASSES OF STRATA IN THE CHOW RING OF R_d

In this section we prove that the subschemes $R_{d,a}^0$ extend to a projective subscheme representing a Chern class and we compute the classes of the closure of the strata $R_{d,a}^0$ in the Quot scheme compactification R_d .

We consider the universal exact sequence over $R_d \times \mathbb{P}^1$,

$$(9) \quad 0 \rightarrow \mathcal{K} \rightarrow \mathcal{O}_{R_d \times \mathbb{P}^1}^4 \rightarrow \mathcal{E} \rightarrow 0.$$

Tensoring (9) with $\pi_2^* \mathcal{O}_{\mathbb{P}^1}(-a-2)$, where π_2 is the projection map over the second factor, we obtain the exact sequence:

$$0 \rightarrow \mathcal{K} \otimes \pi_2^* \mathcal{O}_{\mathbb{P}^1}(-a-2) \rightarrow \mathcal{O}_{\mathbb{P}^1 \times R_d}^4 \otimes \pi_2^* \mathcal{O}_{\mathbb{P}^1}(-a-2) \rightarrow \mathcal{E} \otimes \pi_2^* \mathcal{O}_{\mathbb{P}^1}(-a-2) \rightarrow 0.$$

Let π_1 be the projection map over the first factor. The π_{1*} direct image of the above sequence yields an exact sequence over R_d :

$$\begin{aligned} 0 \rightarrow \pi_{1*}(\mathcal{K} \otimes \pi_2^* \mathcal{O}_{\mathbb{P}^1}(-a-2)) &\rightarrow \pi_{1*}(\mathcal{O}_{R_d \times \mathbb{P}^1}^4 \otimes \pi_2^* \mathcal{O}_{\mathbb{P}^1}(-a-2)) \rightarrow \\ &\rightarrow \pi_{1*}(\mathcal{E} \otimes \pi_2^* \mathcal{O}_{\mathbb{P}^1}(-a-2)) \rightarrow R^1 \pi_{1*}(\mathcal{K} \otimes \pi_2^* \mathcal{O}_{\mathbb{P}^1}(-a-2)) \rightarrow \\ &\rightarrow R^1 \pi_{1*}(\mathcal{O}_{\mathbb{P}^1 \times R_d}^4 \otimes \pi_2^* \mathcal{O}_{\mathbb{P}^1}(-a-2)) \rightarrow R^1 \pi_{1*}(\mathcal{E} \otimes \pi_2^* \mathcal{O}_{\mathbb{P}^1}(-a-2)) \rightarrow 0. \end{aligned}$$

By applying the Serre duality theorem, it follows that

$$(10) \quad R^1 \pi_{1*}(\mathcal{K} \otimes \pi_2^* \mathcal{O}_{\mathbb{P}^1}(-a-2)) \cong \pi_{1*}(\mathcal{K}^\vee \otimes \pi_2^* \mathcal{O}_{\mathbb{P}^1}(a))^\vee$$

and by the base Change theorem, their fibers are isomorphic to

$$H^1(\mathbb{P}^1, \mathcal{K} \otimes \pi_2^* \mathcal{O}_{\mathbb{P}^1}(-a-2)|_{\{q\} \times \mathbb{P}^1}) \cong H^0(\mathbb{P}^1, \mathcal{K}^\vee \otimes \pi_2^* \mathcal{O}_{\mathbb{P}^1}(a)|_{\{q\} \times \mathbb{P}^1})'.$$

In order to see that the sheaf (10) is a bundle, it is enough to check that the cohomology group

$$H^1(\mathbb{P}^1, \mathcal{K}^\vee \otimes \pi_2^* \mathcal{O}_{\mathbb{P}^1}(a)|_{\{q\} \times \mathbb{P}^1}),$$

vanishes or that the group $H^0(\mathbb{P}^1, \mathcal{K} \otimes \pi_2^* \mathcal{O}_{\mathbb{P}^1}(-a-2)|_{\{q\} \times \mathbb{P}^1})$ vanishes, by the Serre duality theorem.

We can assume that $\mathcal{K}|_{\{q\} \times \mathbb{P}^1} \cong \mathcal{O}_{\mathbb{P}^1}(-n) \oplus \mathcal{O}_{\mathbb{P}^1}(n-d)$ with $n \geq 0$ and $n \leq d-n$, therefore $\mathcal{K} \otimes \pi_2^* \mathcal{O}_{\mathbb{P}^1}(-a-2)|_{\{q\} \times \mathbb{P}^1} \cong \mathcal{O}_{\mathbb{P}^1}(-n-a-2) \oplus \mathcal{O}_{\mathbb{P}^1}(n-d-a-2)$. Since the integers $-n-a-2$ and $n-d-a-2$ are both negative, the result follows. For similar reasons, the sheaf

$$(11) \quad R^1 \pi_{1*}(\mathcal{O}_{R_d \times \mathbb{P}^1}^4 \otimes \pi_2^* \mathcal{O}_{\mathbb{P}^1}(-a-2)),$$

is a bundle over R_d .

Remark 4. Since the universal quotient sheaf $\mathcal{E}$ is flat over the R_d Quot scheme, for each $q \in R_d$, $E_q := \mathcal{E}|_{\{q\} \times \mathbb{P}^1}$ is a coherent sheaf over $\mathbb{P}^1$ and therefore it can be written as a direct sum of its locally free part plus its torsion. By the Riemann-Roch theorem, we have that

$$(12) \quad h^0(E_q) - h^1(E_q) = \deg(E_q) + \text{rank}(E_q)(1-g) = d+2.$$

In particular, the degree of E_q is constant for every $q \in R_d$. The sets

$$h^1(E_q) \geq r,$$

where r is an integer, are closed by the Semicontinuity theorem and by the identity (12), they can be also defined as:

$$h^0(E_q) \geq d+2+r.$$

Lemma 4.1. *The Zariski closure $R_{d,a}$ of the stratum $R_{d,a}^0$ in the Quot scheme compactification R_d coincides with the set:*

$$(13) \quad \{q \in R_d \mid h^0(\mathbb{P}^1, E_q \otimes \mathcal{O}_{\mathbb{P}^1}(-a-2)) \geq d+3\}.$$

Proof. First we observe that $R_{d,a}^0 \subset R_{d,a}$ and that every point in the set (13) is in fact in the closure $R_{d,a}$ by Lemma 2.3 and the observation in Remark 4. Now we have to see that every point in the boundary of $R_{d,a}$ belongs to the set (13).

A point $q \in R_d$ is in the boundary, if and only if, E_q has a non-zero torsion T . The contribution of T to the total degree is given by the formula

$$\deg(E_q) = \deg(F) + h^0(T),$$

where F is the locally free part of E_q .

A point q_a in the boundary of $R_{d,a}$ is of the form $\mathcal{O}_{\mathbb{P}^1}^4 \rightarrow \mathcal{O}_{\mathbb{P}^1}(a-a_1) \oplus \mathcal{O}_{\mathbb{P}^1}(b) \oplus \mathcal{O}_{Z_{a_1}}$, or $\mathcal{O}_{\mathbb{P}^1}^4 \rightarrow \mathcal{O}_{\mathbb{P}^1}(a) \oplus \mathcal{O}_{\mathbb{P}^1}(b-b_1) \oplus \mathcal{O}_{Z_{b_1}}$ with a torsion of degree $a_1 > 0$ and $b_1 > 0$ respectively, and $a+b := d$. It obviously satisfies the condition $h^0(\mathbb{P}^1, E_{q_a} \otimes \mathcal{O}_{\mathbb{P}^1}(-a-2)) \geq d+3$, and therefore it belongs to the set (13). $\square$

Theorem 4.2. *$R_{d,a}$ is the locus where the map*

$$(14) \quad R^1 \pi_{1*}(\mathcal{K} \otimes \pi_2^* \mathcal{O}_{\mathbb{P}^1}(-a-2)) \rightarrow R^1 \pi_{1*}(\mathcal{O}_{R_d \times \mathbb{P}^1}^4 \otimes \pi_2^* \mathcal{O}_{\mathbb{P}^1}(-a-2))$$

is not surjective. It is non-empty and has expected codimension $d-2a-1$ as a determinantal variety.

Proof. The map (14) is not surjective in the support of the sheaf

$$(15) \quad R^1 \pi_{1*}(R_d \times \mathbb{P}^1, \mathcal{E} \otimes \pi_2^* \mathcal{O}_{\mathbb{P}^1}(-a-2)),$$

that is, in the points $q \in R_d$ such that

$$h^1(\mathbb{P}^1, E_q \otimes \mathcal{O}_{\mathbb{P}^1}(-a-2)) \geq 1.$$

By the Serre duality theorem and Lemma 4.1, these are the points $q \in R_d$ such that

$$h^0(\mathbb{P}^1, E_q \otimes \mathcal{O}_{\mathbb{P}^1}(-a-2)) \geq d+3.$$

The non-emptiness follows from the fact that given a vector bundle $E \cong \mathcal{O}_{\mathbb{P}^1}(a) \oplus \mathcal{O}_{\mathbb{P}^1}(d-a)$, it is generated by global sections since by assumption $a, d-a \geq 0$ and therefore there exists $f \in R_d^0$ such that $f^*\mathcal{Q} \cong E$.

By applying the Serre duality theorem we get that

$$(16) \quad R^1\pi_{1*}(\mathcal{K} \otimes \pi_2^*\mathcal{O}_{\mathbb{P}^1}(-a-2)) \cong \pi_{1*}(\mathcal{K}^\vee \otimes \pi_2^*\mathcal{O}_{\mathbb{P}^1}(a))^\vee.$$

We apply Riemann-Roch in $\mathbb{P}^1$ in order to compute the dimension of their fibers:

$$\dim H^0(\mathbb{P}^1, \mathcal{K}^\vee \otimes \pi_2^*\mathcal{O}_{\mathbb{P}^1}(a)|_{\{p\} \times \mathbb{P}^1}) = d+2a+2, \quad p \in R_d.$$

As a consequence $\pi_{1*}(\mathcal{K}^\vee \otimes \pi_2^*\mathcal{O}_{\mathbb{P}^1}(a))$ is a bundle of rank $d+2a+2$ over R_d .

Again we see by the Serre duality theorem that

$$(17) \quad R^1\pi_{1*}(\mathcal{O}_{R_d \times \mathbb{P}^1}^4 \otimes \pi_2^*\mathcal{O}(-a-2)) \cong \pi_{1*}(\mathcal{O}_{R_d \times \mathbb{P}^1}^4 \otimes \pi_2^*\mathcal{O}(-a-2))^\vee$$

Since $\mathcal{O}_{\mathbb{P}^1}^4 \otimes \pi_2^*\mathcal{O}_{\mathbb{P}^1}(a)|_{\{p\} \times \mathbb{P}^1}$ is a locally free sheaf over $\mathbb{P}^1$, it can be decomposed over $\mathbb{P}^1$, so that

$$\dim H^0(\mathbb{P}^1, \mathcal{O}_{R_d \times \mathbb{P}^1}^{4\vee} \otimes \pi_2^*\mathcal{O}_{\mathbb{P}^1}(a)|_{\mathbb{P}^1 \times \{p\}}) = 4(a+1).$$

As a consequence $\pi_{1*}(\mathcal{O}_{R_d \times \mathbb{P}^1}^4 \otimes \pi_2^*\mathcal{O}_{\mathbb{P}^1}(a))$ is a trivial bundle of rank $4a+4$ over R_d , that is,

$$\pi_{1*}(\mathcal{O}_{R_d \times \mathbb{P}^1}^4 \otimes \pi_2^*\mathcal{O}_{\mathbb{P}^1}(a)) \cong \mathcal{O}_{R_d}^{4a+4}.$$

Finally, all these facts give us the following morphism of bundles:

$$\pi_{1*}(\mathcal{K}^\vee \otimes \pi_2^*\mathcal{O}_{\mathbb{P}^1}(a)) \xrightarrow{\phi} \mathcal{O}_{R_d}^{4a+4}.$$

The expected codimension of $R_{d,a}$ as a determinantal variety is:

$$((d+2a+2) - (4a+3)) \cdot ((4a+4) - (4a+3)) = d-2a-1 \quad (\S \text{ II.4, [ACGH]}).$$

This codimension coincides with the computed codimension of $R_{d,a}^0$ in 2.2. $\square$

4.0.1. Class of $R_{d,a}$ in $A(R_d)$.

Lemma 4.3. *$R_{d,a}$ is irreducible and does not have components at infinity.*

Proof. Since the subschemes $R_{d,a}$ are the closure of the irreducible sets $R_{d,a}^0$, it follows that they are also irreducible, and consequently cannot have components contained at infinity. $\square$

Proposition 4.4. *If $R_{d,a}$ is either empty or has the expected codimension $d - 2a - 1$, its fundamental class is given by the formula:*

$$[R_{d,a}] = -c_{d-2a-1}(\pi_{1*}(\mathcal{K}^\vee \otimes \pi_2^*(\mathcal{O}_{\mathbb{P}^1}(a)))).$$

Proof. By Lemma 4.3 and Theorem 4.2, the subschemes $R_{d,a}$ are irreducible and have the expected codimension as determinantal varieties. The formula for their Poincare duals is derived from Porteous formula:

$$[R_{d,a}] = \Delta_{d-2a-1,1}(c_t(-\pi_{1*}(\mathcal{K}^\vee \otimes \pi_2^*(\mathcal{O}_{\mathbb{P}^1}(a)))))$$

□

Computation of Chern classes. In order to compute the Chern classes of the bundle $\pi_{1*}(\mathcal{K}^\vee \otimes \pi_2^*(\mathcal{O}_{\mathbb{P}^1}(a)))$ we apply the Grothendieck-Riemann-Roch theorem, so that

$$(18) \quad \text{ch}(\pi_{1*}(\mathcal{K}^\vee \otimes \pi_2^*(\mathcal{O}_{\mathbb{P}^1}(a)))) = \pi_{1*}(\text{Td}(R_d \times \mathbb{P}^1 / R_d) \cdot \text{ch}(\mathcal{K}^\vee \otimes \pi_2^*(\mathcal{O}_{\mathbb{P}^1}(a)))).$$

$$(19) \quad c_i(\mathcal{K}) = t_i + h u_{i-1} \quad (1 \leq i \leq 2),$$

$t_i, u_i \in A^i(R_d), i = 1, \dots, n-1$ as in [Str], in particular $u_0 = -d$.

First we compute the Chern classes of $\mathcal{K}^\vee \otimes \pi_2^*(\mathcal{O}_{\mathbb{P}^1}(a))$:

$$\begin{aligned} c_1(\mathcal{K}^\vee \otimes \pi_2^*(\mathcal{O}_{\mathbb{P}^1}(a))) &= c_1(\mathcal{K}^\vee) + c_1(\pi_2^*(\mathcal{O}_{\mathbb{P}^1}(a))) \cdot c_0(\mathcal{K}^\vee) \\ &= h(2a + d) - t_1, \end{aligned}$$

$$\begin{aligned} c_2(\mathcal{K}^\vee \otimes \pi_2^*(\mathcal{O}_{\mathbb{P}^1}(a))) &= c_2(\mathcal{K}^\vee) + c_1(\mathcal{K}^\vee) \cdot c_1(\pi_2^*(\mathcal{O}_{\mathbb{P}^1}(a))) + c_1(\pi_2^*(\mathcal{O}_{\mathbb{P}^1}(a)))^2 \cdot c_0(\mathcal{K}^\vee) \\ &= -a t_1 h - t_2 - h u_1, \end{aligned}$$

$$c_3(\mathcal{K}^\vee \otimes \pi_2^*(\mathcal{O}_{\mathbb{P}^1}(a))) = a h t_2.$$

Let us denote by ch_i , the i -homogeneous part of the Chern character of a bundle.

$$\text{ch}_0(\mathcal{K}^\vee \otimes \pi_2^*(\mathcal{O}_{\mathbb{P}^1}(a))) = 2,$$

$$\text{ch}_1(\mathcal{K}^\vee \otimes \pi_2^*(\mathcal{O}_{\mathbb{P}^1}(a))) = -t_1 + h(d + 2a),$$

$$\text{ch}_2(\mathcal{K}^\vee \otimes \pi_2^*(\mathcal{O}_{\mathbb{P}^1}(a))) = \frac{1}{2}[t_1^2 - 2t_1 h(2a + d) + 2at_1 h + 2t_2 + 2hu_1],$$

$$\text{ch}_3(\mathcal{K}^\vee \otimes \pi_2^*(\mathcal{O}_{\mathbb{P}^1}(a))) = -\frac{1}{6}[t_1^3 - 3(d + 2a)h t_1^2 + 3t_1 t_2 + 3hu_1 t_1 - 3a h t_1^2 -$$

$$-3u_2 h - 3a h t_2],$$

Lemma 4.5.

$$\text{ch}_n(\mathcal{K}^\vee \otimes \pi_2^*(\mathcal{O}_{\mathbb{P}^1}(a))) = \text{coeff}_{(t^n)} \left(\sum_n \frac{(-1)^{n-1}}{n} \sum_j c_j(\mathcal{K}^\vee \otimes \pi_2^*(\mathcal{O}_{\mathbb{P}^1}(a))) t^j \right)^n.$$

Proof. The Chern character of a bundle E is defined by the formula:

$$\text{ch}(E) = \sum_i e^{\alpha_i} = \sum_i \sum_n \frac{(x_i)^n}{n!} \in H^*(X, \mathbb{Q}).$$

If we call σ_i to the symmetric functions, that is,

$$\sum_{r=0}^n \sigma_r t^r = \prod_{i=1}^n (1 + x_i t),$$

we get that

$$\begin{aligned} \log(1 + \sigma_1 t + \sigma_2 t^2 + \dots) &= \log \prod_i (1 + x_i t) \\ &= \sum_i \sum_n \pi_2^* \frac{(-1)^{n-1}}{n} (x_i t)^n = \sum_n \frac{(-1)^{n-1}}{n} (\sum x_i^n t^n). \end{aligned}$$

and from here the formula above follows. $\square$

Now applying (18), we have

$$\text{ch}(\pi_{1*}(\mathcal{K}^\vee \otimes \pi_2^* \mathcal{O}_{\mathbb{P}^1}(a))) = \pi_{1*}((1 + h)(\text{ch}(\mathcal{K}^\vee \otimes \mathcal{O}_{\mathbb{P}^1}(a)))) .$$

Therefore the Chern classes of $\pi_{1*}(\mathcal{K}^\vee \otimes \pi_2^* \mathcal{O}_{\mathbb{P}^1}(a))$ are:

$$\begin{aligned} c_1(\pi_{1*}(\mathcal{K}^\vee \otimes \pi_2^* \mathcal{O}_{\mathbb{P}^1}(a))) &= -t_1(a + d + 1) + u_1, \\ c_2(\pi_{1*}(\mathcal{K}^\vee \otimes \pi_2^* \mathcal{O}_{\mathbb{P}^1}(a))) &= \frac{1}{2} c_1^2 - \left[\frac{1}{2} (2a + d) t_1^2 - \frac{1}{2} a t_1^2 + \frac{1}{2} (2a + d) t_2 - \frac{1}{2} u_1 t_1 \right], \\ &\vdots \end{aligned}$$

Lemma 4.6.

$$\begin{aligned} c_n(\pi_{1*}(\mathcal{K}^\vee \otimes \pi_2^* \mathcal{O}_{\mathbb{P}^1}(a))) &= \\ - \sum_{r=1}^n \frac{(-1)^{r-1}}{n} r! \text{ch}_r(\pi_{1*}(\mathcal{K}^\vee \otimes \pi_2^* \mathcal{O}_{\mathbb{P}^1}(a))) c_{n-r}(\pi_{1*}(\mathcal{K}^\vee \otimes \pi_2^* \mathcal{O}_{\mathbb{P}^1}(a))). \end{aligned}$$

Proof. For each $r \geq 1$ the r 'th power sum is:

$$p_r = \sum x_i^r = m_{(r)}.$$

The generating function for the p_r is:

$$(20) \quad p(t) = \sum_{r \geq 1} p_r t^{r-1} = \sum_{i \geq 1} \sum_{r \geq 1} x_i^r t^{r-1} = \sum_{i \geq 1} \frac{d}{dt} \log \frac{1}{1 - x_i t}.$$

$$P(t) = \frac{d}{dt} \prod_{i \geq 1} (1 - x_i t)^{-1} = \frac{d}{dt} \log H(t) = \frac{H'(t)}{H(t)}$$

$$(21) \quad P(-t) = \frac{d}{dt} \log E(t) = \frac{E'(t)}{E(t)}.$$

From (20) and (4.0.1) we get that

$$n \sigma_n = \sum_{r=1}^n (-1)^{r-1} p_r \sigma_{n-r},$$

and the formula above follows. $\square$

5. ANOTHER COMPACTIFICATION OF THE SPACE R_d^0 .

Now we will consider the coarse moduli space $\overline{M}_{0,n}(G(2,4),d)$ of degree d Kontsevich stable maps from n -pointed, genus 0 curves to $G(2,4)$. It is smooth, has dimension $4d + 1 + n$ and there is a universal family over it:

$$\begin{array}{ccc}
 0 \rightarrow \mathcal{N} \rightarrow \mathcal{O}^4 \rightarrow \mathcal{Q} \rightarrow 0 \\
 \downarrow \\
 \overline{M}_{0,n+1}(G(2,4),d) \xrightarrow{e_i} G(2,4) \\
 \pi_i \downarrow \uparrow \sigma_i \\
 \overline{M}_{0,n}(G(2,4),d)
 \end{array}$$

Here $\pi_n : \overline{M}_{0,n+1}(G(2,4),d) \rightarrow \overline{M}_{0,n}(G(2,4),d)$ is the forgetful morphism, which consists of forgetting the mark p_1 and σ_i is the section corresponding to the mark p_i . The image of σ_i is the (closure of the) locus of maps whose source curve has two components, one of which is rational, carries just the two marks p_i and p_{n+1} and is contracted by μ . For $n = 3$, the space $\overline{M} := \overline{M}_{0,3}(G(2,4),d)$ provides a compactification of the space R_d^0 . The marked points yield canonical line bundles $\mathcal{L}_i = e_i^*(T_1)$ on $\overline{M}$ via the i -evaluation map, where T_1 is the hyperplane class in $G(2,4)$. Since the boundary is of pure codimension 1 in $\overline{M}_{0,n}(G(2,4),d)$, each irreducible component is a Weil divisor. Let Δ be the set of components of the boundary. For $d \geq 1$, Weil divisors are obtained on $\overline{M}_{0,n}(G(2,4),d)$ by considering the locus corresponding to maps meeting an a -plane and a b -plane on the Grassmannian. We denote the corresponding classes in $\text{Pic}(\overline{M}) \otimes \mathbb{Q}$ by $\mathcal{A}$ and $\mathcal{B}$ respectively. Let w_π denote the relative dualizing sheaf, then a map $\mu \in \overline{M}_{0,n}(G(2,4),d)$ is stable if and only if $w_\pi(p_1 + \dots + p_n) \otimes \mu^*(T_3)$ is ample. In particular $\deg(w_\pi(p_1 + \dots + p_n) \otimes \mu^*(T_3)) > 0$, so that if $E \subset C$ such that $\dim \mu(E) = 0$, then $\deg(w_\pi(p_1 + \dots + p_n)) > 0$.

Theorem 5.1. *If $d \geq 1$, then $\text{Pic}(\overline{M})$ is generated by $\Delta \cup \{\mathcal{A}\} \cup \{\mathcal{L}_i\}_{i=1,2,3}$.*

Proof. Consider the universal family of stable degree d maps of 3-pointed curves $R_d^{00} \times \mathbb{P}^1 \rightarrow G(2,4)$. By the universal property, there is an injection

$$R_d^{00} \hookrightarrow \overline{N}.$$

The complement of R_d^{00} is the boundary of $\overline{N}$, that is, the locus corresponding to maps with reducible domains. Since the complement of R_d^{00} in R_d^0 is of codimension greater or equal to 2 (see [Mar]), the Picard group of R_d^{00} coincides with the Picard group of R_d^0 . The divisor $\mathcal{A}$ restricted to R_d^0 , i.e. to the moduli points corresponding to stable maps with irreducible source, is linearly equivalent to D_i and $\mathcal{L}_i$ is linearly equivalent to the corresponding divisor Y_i . In these points, markings are not necessary since the maps are already stable. In section 3, we argued that the divisors Y_i and D_i constitute a basis of $A^1(R_d^0)$, so that $A_{4d+3}(\overline{M})$ is generated by the boundary Δ and the Weil divisors $\mathcal{L}_i$ ($i = 1, 2, 3$), and $\mathcal{A}$. We note that since $Y_1 \sim Y_2 \sim Y_3$ in R_d^0 , each one of the $\mathcal{L}_i$ generates the Picard group. $\square$

If $d = 0$, then $\overline{M}_{0,n}(G(2,4), 0) \cong \overline{M}_{0,n} \times G(2,4)$ and $\mathcal{L}_i$ is the pull-back of $\mathcal{O}_{G(2,4)}(1)$ from the second factor.

Boundary of $\overline{M}_{0,n}(G(2,4), d)$. The irreducible components of the boundary are in bijective correspondence with the data of weighted partitions $(A \cup B, d_A, d_B)$, where

- $A \cup B$ is a partition of the set of marked points. There are only 2 possible partitions: $1+2, 3+0$.
- $d_A + d_B = d, d_A \geq 0, d_B \geq 0$,
- if $d_A = 0$ (resp. d_B), then $|A| \geq 2$ (resp. $|B| \geq 2$).

A component $(A \cup B, d_A, d_B)$ corresponds generically to a map with a union of $\mathbb{P}^1$'s as its domain. Each one gives a curve in $G(2,4)$ of degrees d_A and d_B , respectively.

5.1. Stratification of $\overline{M}$. The stratification of the space R_d^0 given in 2.5 can be extended to $\overline{M}$, we can even compute the classes of the strata in the Chow ring $A(\overline{M})$, but they have no enumerative meaning.

5.1.1. Closure of the stratum in $\overline{M}$. The closure of the stratum has already been studied by I. Coskun in [Cos] by taking flat limits of surfaces in one parameter families.

Let $V_{d,a}$ denote the closure of $R_{d,a}^0$ in $\overline{M}$. Let K^i be the boundary component corresponding to the degree partition $i + (d - i) = d$. Geometrically these boundary points correspond to reducible curves in the Grassmannian $G(2,4)$ of degree $i, d - i$ respectively, or equivalently a union of irreducible rational ruled surfaces of degree i and $d - i$ in $\mathbb{P}^3$.

A generic intersection point of the boundary with $V_{d,a}$ corresponds to a moduli point,

$[C, p_1, p_2, p_3, \mu : C \rightarrow G(2,4)]$, where $C = C_1 \cup C_2$ is a union of two $\mathbb{P}^1$'s with $\mu^* \mathcal{Q}|_{C_1} \cong \mathcal{O}_{\mathbb{P}^1}(a_1) \oplus \mathcal{O}_{\mathbb{P}^1}(b_1)$, $a_1 + b_1 = i$ and $\mu^* \mathcal{Q}|_{C_2} \cong \mathcal{O}_{\mathbb{P}^1}(a_2) \oplus \mathcal{O}_{\mathbb{P}^1}(b_2)$, $a_2 + b_2 = d - i$. The integers a_j, b_j satisfy $a_j \geq 0, b_j \geq 0, \sum_j (a_j + b_j) = d$ and $a_1 + a_2 \leq a$. This last condition means that the flat limit of the directrices is a connected curve of degree a whose restriction to each of the surfaces is in a section class. Conversely, given any connected curve C of degree $a \leq \sum (a_j + b_j)/2$ whose restriction to each component is in a section class, there exists a one parameter family of $X_{a,b}$ specializing to the reducible surface such that the limit of the directrices is C .

The stratification given in section 2 is extended to $\overline{M}$ by defining the stratum as

$$(22) \quad \overline{M}_{d,a} := \{\mu \in \overline{M} \mid h^1(\mu^* \mathcal{Q} \otimes w_\pi(-a p_i)) \geq 1\}.$$

This set is a closed subset of $\overline{M}$ by the Semicontinuity Theorem.

Proposition 5.2. $\overline{M}_{d,a}$ is the locus where the map

$$(23) \quad R^1 \pi_{i*}(e_i^*(\mathcal{N}) \otimes w_\pi(-a p_i)) \rightarrow R^1 \pi_{i*}(e_i^*(\mathcal{O}^4) \otimes w_\pi(-a p_i))$$

is not surjective.

Proof. We proceed as in section 4, considering the long exact sequence:

$$\begin{aligned}
0 \rightarrow \pi_{i*}(e_i^* \mathcal{N} \otimes w_\pi(-a p_i)) &\rightarrow \pi_{i*}(e_i^* \mathcal{O}^4 \otimes w_\pi(-a p_i)) \rightarrow \pi_*(e_i^* \mathcal{Q} \otimes w_\pi(-a p_i)) \\
&\rightarrow R^1 \pi_*(e_i^* (\mathcal{N}) \otimes w_\pi(-a p_i)) \rightarrow R^1 \pi_*(e_i^* (\mathcal{O}^4) \otimes w_\pi(-a p_i)) \rightarrow \\
&R^1 \pi_*(e_i^* (\mathcal{Q}) \otimes w_\pi(-a p_i)) \rightarrow 0.
\end{aligned}$$

The map (23) is not surjective in the support of the sheaf $R^1 \pi_{1*}(e_i^* (\mathcal{Q}) \otimes w_\pi(-a p_i))$, that is at the moduli points $[C, p_1, p_2, p_3, \mu : C \rightarrow G(2, 4)]$ where $h^1(C, \mu^* \mathcal{Q} \otimes w_\pi(-a p_i)) \geq 1$, and these are the points defining the stratum (22). $\square$

5.1.2. *Enumerative geometry of $\overline{M}_{d,a}$.* We would like to know which points in the boundary of $\overline{M}$ satisfy the condition:

$$h^1(C, \mu^* \mathcal{Q} \otimes w_\pi(-a p_i)) \geq 1.$$

The next theorem will address this question.

Recalling the notation of [Mar], Q_d will be the number of rational ruled surfaces through $4d + 1$ -points in $\mathbb{P}^3$. The numbers Q_d have the following geometric interpretation in terms of $\mathbb{Q}$ -divisors. The number Q_d is the zero-cycle $\mathcal{A}^{4d+1}$ in the Chow ring of $\overline{M}_{0,0}(G(2, 4), d)$. The number Q_d^b will denote the zero-cycle $\mathcal{A}^{4d} \mathcal{B}$, or equivalently the Gromov-Witten invariant, $I_{0,4d+1,d}(T_a, \overset{4d}{\dots}, T_a, T_b)$. The following table lists the numbers Q_d^b for each degree $3 \leq d \leq 9$. These numbers correspond geometrically to rational ruled surfaces of degree d through $4d$ points in $\mathbb{P}^3$ and tangent to a fixed plane.

d	Q_d^b	$4d$
1	0	4
2	2	8
3	1824	12
4	3094440	16
5	15383867920	20
6	188115939619440	24
7	3000224401806629008	28
8	219761533783440334862592	32
9	17394248462381072210049044320	36

Theorem 5.3. $V_{d,a} \subset \overline{M}_{d,a}$ and $V_{d,a} = \overline{M}_{d,a}$ only when $a = \frac{d}{2} - 1$. In this case, the stratum $R_{d,a}^0$ is of pure codimension 1 and its degree is given by the formula

$$\frac{1}{2} d^3 \cdot (Q_d + Q_d^b).$$

Proof. By semicontinuity (see [Har], III. 12.8), the points $[C, p_1, p_2, p_3, \mu : C \rightarrow G(2, 4)]$ in the closure of $R_{d,a}^0$ in $\overline{M}$ satisfy the condition 5.1.2. The converse is false. We will construct some families satisfying the condition but not in the closure of the stratum.

Construction. Grothendieck duality implies

$$h^1(C, \mu^* \mathcal{Q} \otimes w_\pi(-a p_i)) = h^0(C, \mu^* \mathcal{Q}^\vee \otimes \mathcal{O}(ap_i)).$$

For a general boundary point, the source curve $C = C_1 \cup C_2$ is a nodal curve. There is a canonical exact sequence:

$$0 \rightarrow \mu^* \mathcal{Q}^\vee|_C \otimes \mathcal{O}(ap_i) \rightarrow \mu^* \mathcal{Q}^\vee|_{C_1} \otimes \mathcal{O}(ap_i) \oplus \mu^* \mathcal{Q}^\vee|_{C_2} \otimes \mathcal{O}(ap_i) \rightarrow \mathbb{C}^2 \rightarrow 0.$$

This gives the following injections:

$$\begin{array}{c} H^0(C_1, \mu^* \mathcal{Q}^\vee|_{C_1} \otimes \mathcal{O}(ap_i)) \rightarrow H^0(C, \mu^* \mathcal{Q}^\vee|_C \otimes \mathcal{O}(ap_i)) \\ \uparrow \\ H^0(C_2, \mu^* \mathcal{Q}^\vee|_{C_2} \otimes \mathcal{O}(ap_i)). \end{array}$$

Since $C_i \cong \mathbb{P}^1$, it can be assumed that $\mu^* \mathcal{Q}|_{C_1} \cong \mathcal{O}(a_1) \oplus \mathcal{O}(b_1)$ and $\mu^* \mathcal{Q}|_{C_2} \cong \mathcal{O}(a_2) \oplus \mathcal{O}(b_2)$ with a_1, a_2, b_1, b_2 being integers greater than or equal to 0 and satisfying $\sum_i a_i + b_i \leq d$. Also we assume that $P_i \in C_1$, so that $h^0(C_2, \mu^* \mathcal{Q}^\vee|_{C_2} \otimes \mathcal{O}(ap_i)) = h^0(C_2, \mathcal{O}(-a_2) \oplus \mathcal{O}(-b_2))$, and $h^0(C_1, \mu^* \mathcal{Q}^\vee|_{C_1} \otimes \mathcal{O}(ap_i)) = h^0(C_1, \mathcal{O}(a - a_1) \oplus \mathcal{O}(a - b_1))$.

- (1) If $a < a_1$ and $a = b_1$, condition 5.1.2 is still satisfied, but these points are not in the closure $V_{d,a}$, (see 5.1.1).
- (2) If $a = a_1$ and $a_2 = 0$, then there exists a family of surfaces degenerating to a union of a cone of degree b_2 and a scroll X_{a_1, b_1} , so that these points are in the closure.
- (3) If $a = a_1$ and $a_2 > 0$, then by (5.1.1) the union $X_{a_1, b_1} \cup X_{a_2, b_2}$ cannot be a flat limit of scrolls.

The degree of the stratum $\overline{M}_{\frac{d}{2}-1}$. Note that by degree we mean, the number of rational ruled surfaces of this specifying type through $4d+1$ points, which we studied in [Mar].

For $a = \frac{d}{2} - 1$, proposition 2.2 shows that R_a^0 is a one codimensional subvariety. Its class $(a - d + 1)Y + D$ in the Chow ring $A(R_d^0)$, is given by Porteous formula in section 2 and Lemma 3.2. Because the divisors in the Picard group $A^1(\overline{N})$ intersect the boundary transversally, it follows by (5.1) that this divisor extends in the moduli compactification $\overline{M}$, to the divisor $(a + d - 1)\mathcal{L}_i + \mathcal{A}$. Therefore, the degree of the stratum $\overline{M}_{\frac{d}{2}-1}$, that is, the degree of the variety parametrizing rational ruled surfaces with a minimal directrix of degree $\frac{d}{2} - 1$, is the degree of the top codimensional class $((a + d - 1)\mathcal{L}_i + \mathcal{A}) \cdot \mathcal{A}^{4d} \cdot \mathcal{L}_1 \cdot \mathcal{L}_2 \cdot \mathcal{L}_3$ on $\overline{M}$, determined by general a -planes $T_{a_1}, \dots, T_{a_{4d}}$ and hyperplanes T_{3_j} , $j = 1, 2, 3$, (see [Mar]). Now since $\mathcal{L}_1^2 \cdot \mathcal{A}^{4d} = Q_d + Q_d^b$ in the Chow ring $A(\overline{M}_{0,1}(\mathbb{G}(2,4), d))$, the result follows. $\square$

REFERENCES

- [ACGH] E. Arbarello, M. Cornalba, P.A. Griffiths, J. Harris. *Geometry of Algebraic Curves* (Volume I).
- [Har] R. Hartshorne. *Algebraic Geometry*, GTM 52, Springer, 1977.
- [Ber1] A. Bertram, *Quantum Schubert Calculus*, Adv. Math. **128** (1997), no. 2, 289-305.

- [Ber2] A. Bertram, *Towards a Schubert Calculus for Maps from a Riemann Surface to a Grassmannian*, Internat. J. Math. **5** (1994), no.6, 811-825.
- [Cos] I. Coskun, *Degenerations of surface scrolls and the Gromov Witten invariants of Grassmannians*, Journal of Algebraic Geometry, **15** (2006), p. 223-284.
- [FP] W. Fulton and R. Pandharipande, *Notes on stable maps and quantum cohomology-Santa Cruz, 1995, 45-96*, Proc. Sympos. Pure Math. **62**.
- [Mar] C. Martínez, *The degree of the variety of rational ruled surfaces and Gromov-Witten invariants*, Trans. Amer. Math. Soc. 358 (2006), 11-24.
- [Mar2] C. Martínez, *On the cohomology of Brill Noether loci over Quot schemes*, math.AG/0602491.
- [Pan1] R. Pandharipande, *Intersection of Q -divisors on Kontsevich's moduli space $\overline{M}_{0,n}(\mathbb{P}^r, d)$ and Enumerative Geometry*, Transactions of the American Mathematical Society, no.4, April 1999, pages 1481-1505.
- [Pan2] R. Pandharipande, *The Canonical Class of $\overline{M}_{0,n}(\mathbb{P}^r, d)$ and Enumerative Geometry*, Internacional Mathematics Research Notices, no. 4, (1997), 173-186.
- [RRW] M.S. Ravi, J. Rosenthal, X.Wang, *Degree of the generalized Plücker embedding of a quot scheme and quantum cohomology*, Math. Ann. 311 (1998), no. 1, 11-26.
- [Sha] S. S. Shatz, *The decomposition and specialization of algebraic families of vector bundles* Compositio Mathematica, 1977.
- [Str] S. A. Strømme. *On parametrized rational curves in Grassmann Varieties*, 251-272 Lecture Notes in Math, 1266, Springer, Berlin, 1987.
- [Vak] R. Vakil, *The Enumerative Geometry of rational and elliptic curves in Projective space*, J. Reine Angew. Math. **529** (2000), 101-153.

INSTITUT FOR MATEMATIKE FAG (CTQM), AARHUS UNIVERSITET, NY MUNKEGADE,
8000 AARHUS, DENMARK

E-mail address: cmartine@imf.au.dk